



Points to note Before tooth removal

**Before tooth removal,
Kindly Inform your Doctor If you are ,...**

- 1 Suffering from any illness**
- 2 Taking any medicines**
- 3 Allergic to any medicines**



Dental extraction is almost a painless procedure as it is done under local anaesthesia.

If any pain is felt during the procedure, it may be because the tooth is infected at the root tip. Thus, the procedure can be continued only after taking medications prescribed by the doctor.

**Proper food should be
taken before tooth
removal .**

**If your Doctor has
prescribed any medicines
to be taken
Before tooth removal,
kindly do so.**

**Pregnant ladies and breast
feeding mothers kindly
inform the fact to the
Doctor.**

YOUR CLINIC DETAILS WILL BE ADDED HERE

Care after Tooth removal

1



Bite gently, but firmly on the COTTON GAUZE. This will control bleeding. Remove it after the bleeding has stopped.

2



Placing an ICE-PACK On the cheek will reduce bleeding and swelling.

3



AVOID HOT, STICKY & HARD FOOD.
Do not use STRAW for drinking.

4



Take only SOFT, COLD FOOD

5



Gargle mildly with LUKE WARM SALT WATER 8 hours after tooth removal. Do so 3 times daily for 7 days.

6



AVOID HARD WORK & SPORTS for some days.

7



NO SMOKING , NO ALCOHOL After tooth removal and during medication.

8



Take the prescribed MEDICINES at the right time.

9



If there is a SUTURE at the site of tooth removal, your Doctor will remove it at the next appointment.

consult your Doctor if there is ,...

Severe pain, after the day of tooth removal.

Uncontrolled bleeding or swelling.

Rashes , itching ,swelling or redness of the eyes , after taking medicines.

YOUR CLINIC DETAILS WILL BE ADDED HERE



Impacted Teeth

In this condition, the majority of the tooth portion remains embedded (impacted) in the jaw bone.

Removal of this tooth, requires much care & effort than a normal tooth removal.

Which teeth are most often Impacted ?

Wisdom tooth or Third molar



They are the most commonest impacted tooth.

Maxillary (upper jaw) Canine tooth



The second most common teeth to be impacted .

YOUR CLINIC DETAILS WILL BE ADDED HERE

Impacted Teeth

Symptoms of Impacted teeth

Red, swollen, or bleeding gums

Bad breath

Bad taste in your mouth

Difficulty in opening your mouth

**Pain when opening your mouth,
or when chewing and biting.**

YOUR CLINIC DETAILS WILL BE ADDED HERE


Orthognathic Surgery

for correction of facial deformities

Orthognathic surgery, or corrective jaw surgery is performed by an oral and maxillofacial surgeon.

The aim is to correct, ... skeletal and dental irregularities, including the misalignment of jaws and teeth.

Orthognathic surgery, improves the overall aesthetic appearance and functions like chewing, speaking and breathing.



Before

After



Before

After



Before

After

YOUR CLINIC DETAILS WILL BE ADDED HERE



Cleft

lip and palate

How does a baby get a cleft lip/palate?

Cleft lip and cleft palate occur very early in pregnancy, while the baby is developing inside the mother.

Clefting results when there is not enough tissue in the mouth or lip area, and the remaining tissue does not join together.

Complications depending on the type and severity of the cleft.

- Difficulty in feeding.
- Ear infections and hearing loss.
- Dental problems.
- Speech difficulties.
- Emotional and behavioral problems due to differences in physical appearance

Factors that are responsible for the occurrence of a cleft lip/palate.

- Parents with a family history of cleft lip or palate face a higher risk of having a baby with a cleft.
- In pregnant women who smoke cigarettes, drink alcohol or take certain medications.
- Having diabetes during pregnancy .
- Being obese during pregnancy.

WATER MARK . WATER MARK
www.smartdocposters.com

WATER MARK . WATER MARK

YOUR CLINIC DETAILS WILL BE ADDED HERE



Cleft

lip and palate

Cleft lip and palate can be surgically corrected!

- The aim of surgical treatment is to improve the child's ability to eat, speak and hear normally and to achieve a normal facial appearance.
- Surgery often involves a team of Doctors and experts .
- Surgeries are performed in this order:
Within the first 12 months of age - Cleft lip repair is done.
By the age of 18 months -Cleft palate repair is done.
Between age 2 and late teen years -Follow-up surgeries are done.
- The surgery takes place in a hospital and is done under general anesthesia, so the child won't feel any pain during or after surgery.



YOUR CLINIC DETAILS WILL BE ADDED HERE